

DIFFERENCES IN CHEMICAL COMPOSITION AMONG PROVENANCES OF TREE FODDER SPECIES IN A SUBHUMID ENVIRONMENT: RELATION TO USE AS SUPPLEMENTS

B.H. Dzowela¹, L. Hove² and P.L. Mafongoya³

¹ SADC-ICRAF Agroforestry Project, P.O. Box CY 594, Causeway, Harare,

² Makoholi Research Station, P/Bag 9248, Masvingo, Zimbabwe

³ Agronomy Institute, Department of Research and Specialist Services, P.O. Box CY550, Causeway, Harare, Zimbabwe

ABSTRACT

Studies were initiated in a subhumid Southern African environment to assess the chemical composition and nutritive value of *Acacia angustissima* (Mill.) Kuntze, *Calliandra calothyrsus* Meiss. and *Leucaena* species. The objective was to determine variability in crude protein (CP) concentration, acid detergent fibre (ADF), neutral detergent fibre (NDF), and soluble and insoluble proanthocyanidins from fodder samples of species, subspecies and provenances. A wide range was found in these parameters. The implications of these chemical composition factors, especially proanthocyanidins, on the use of these browse fodders in livestock feeding systems are discussed.

KEYWORDS

Tree fodders, crude protein, fibres, proanthocyanidins, digestibility, supplements, feeding systems.

INTRODUCTION

Leguminous trees produce high quality fodder. However, their major constraint is their chemical composition and anti-nutritional factors, and how these characteristics affect their use in feeding systems. Factors contributing to high quality forage, are high concentrations of N and low concentrations of lignin and polyphenolic secondary plant compounds. Thus, the chemical composition of fodder from tree fodders could have a strong influence on its use (Ndlovu *et al.*, 1997). The present study was undertaken to assess the levels of chemical composition variability in fodder tree species and provenances so as to relate this variability to potential use as feeds.

MATERIALS AND METHODS

Three evaluation studies involving (1) *Leucaena* genus, (2) *Calliandra calothyrsus* Meissen. and (3) *Acacia angustissima* (Mill.) Kuntze. were carried out at Domboshawa Agroforestry Research site in subhumid Zimbabwe. In the first experiment, the different species and provenances were planted out in fodderbanks. Forage (leaf and petioles) were analysed for the concentration of acid detergent fibre content (ADF), neutral detergent fibre (NDF) (Robertson and van Soest, 1981) and crude protein (AOAC, 1990). The content of proanthocyanidins (condensed tannins) was assayed by the butanol-HCl method of Porter *et al.* (1986) while the soluble tannins were determined by the method of Reed *et al.* (1985). *In vitro* dry matter digestibility (IVDMD) was determined by the Tilley and Terry method (1963). In the second experiment, dry matter (DM) and nitrogen rumen degradations after 24 and 48 hours of incubation in the rumen, were measured using the nylon bag technique (Orskov *et al.*, 1980). In addition, the water soluble DM and N-fractions of the forages were determined. Nutrient intake, digestion and N-balance were measured in the 3rd experiment using sixteen 28-month old goats (average weight 26 ± 4.2 kg). The digestion trial involving 4 goats per treatment, was conducted over a 21-day period which consisted of 14-day adaptation and 7-day data collection periods. The goats were fed natural grass hay *ad libitum* and all three *Leucaena* forages were offered at the rate of 120 g/hd/day.

RESULTS

a) ***Leucaena* species and provenances evaluation:** The comparison

of nitrogen contents between *L. leucocephala* cv. Cunningham (LL), *L. diversifolia* subspecies *stenocarpa* cv. OFI 53/88 (LD) and *L. pallida* (LP) cv. of CPI 58980 (syn. *L. esculenta* subspecies *paniculata*), showed the contents to be high (Table 1). However, 50% or more of this N was bound to NDF. All three species contained substantial amounts of insoluble proanthocyanidins (PAS) which were much higher than that of veld hay. The chemical composition of other *Leucaena* species showed a great deal of variability in crude protein level (Table 2); range of low values was around 17.0% in *L. diversifolia* subsp. *diversifolia* and *stenocarpa* to the high in *L. leucocephala* (27.6%), *L. salvadorensis* (25.6%) and *L. shannonii* subsp. *magnifica* (24.5%). Among the species with low levels of NDF were *L. leucocephala* hybrids with *L. diversifolia*, *L. diversifolia* subsp. *stenocarpa* and *diversifolia* with NDF values below 40%. Among the highest were *L. shannonii* subsp. *magnifica*, *L. pallida* and one *L. diversifolia* provenance subsp. *diversifolia* (OFI 45/87). Similarly, a great deal of variability was observed in PAS concentration between species and subspecies and provenances. The highest values were in *L. pallida* and the two *L. diversifolia* subsp. *stenocarpa* provenances (OFI 35/88 and OFI 53/88). When three contrasting *Leucaena* spp. (*L. diversifolia*, *L. pallida* and *L. leucocephala*) were fed as supplements to low quality native grass hay, there was an increase in DM intake while intake of digestible DM was only increased by supplementing with *L. pallida* and *L. leucocephala*. However, there were no differences ($P > 0.05$) between treatments in terms of digestible NDF intake. Excretion of N was mainly through faeces (over 90 percent) for all diets. Feeding native grass hay alone or with *L. pallida* or *L. diversifolia* resulted in a negative N-balance. Animals fed *L. diversifolia* had lower ($P < 0.05$) urine N relative to the other animals. This is indicative of the protection from rumen microbes of N by proanthocyanidins in *L. diversifolia*. This result is consistent with rumen degradation results (Table 1).

b.) ***Acacia angustissima* provenances evaluation:** A wide variability was observed in the parameters considered (CP, ADF, NDF, soluble and insoluble PAS and IVDMD). Generally, all provenances had higher CP contents in the leaf than in the stem because the leaf does not contain as much structural material as the stem (data not shown). The range in the leaf IVDMD was from 42% in OFI 65/92 to only 50.9% in OFI 34/88 which was similar to the digestibility coefficients recorded in OFI 66/92 (50.4%) and OFI 68/92 (50.7%). The provenances with high digestibility in *A. angustissima* were superior to *C. calothyrsus* (41% IVDMD) and not very much lower than *L. pallida* (IVDMD of 53.1%).

c) ***Calliandra calothyrsus* provenances evaluation:** A wide variability was found in the parameters measured (ADF, NDF, insoluble and soluble PAS and IVDMD). Some provenances, notably OFI 12/91, OFI 61/92 and to a lesser degree OFI 62/92 appear to have digestibility values that are at least 15% higher than the control provenance, OFI 9/89.

DISCUSSION AND CONCLUSION

From the data on *Leucaena* spp. it is apparent that 3 groups of *Leucaena* are available. The first group is one of low NDF, low to

medium PAS and high crude protein concentrations as represented by *L. leucocephala*, its two hybrids with *L. diversifolia*, and *L. shannonii* and *L. salvadorensis*. They represent materials of high nutritive value. Then there is the group which is represented by *L. pallida* and *L. diversifolia* subspecies *stenocarpa* and *diversifolia* to some extent. This group is characterized by high concentrations of PAS with an ABU. value in excess of 45, CP concentration below 20% and NDF values above 35%. These are materials of low nutritive value. The third group is an intermediate group to which the rest of the species, subspecies and provenances belong (including most of the *L. esculenta* species and subspecies). The differences in the world-acclaimed resistance in *L. esculenta* (syn. *L. pallida*) and *L. diversifolia* to the psyllid pest could be related to their condensed tannins content. The efficiency with which these diverse *Leucaena* species and provenances will fit into feeding systems will vary also. Our work has shown lower IVDMD values for *L. diversifolia* and *L. pallida* in addition to higher concentrations of condensed tannin than in *L. leucocephala*. These differences resulted in negative N-balance when the *L. diversifolia* and *L. pallida* were fed as supplements to poor quality native grass hay. These data suggest that consumption of the two *Leucaena* forages (*diversifolia* and *pallida*) may be limited by the low digestibility/degradability especially for *L. diversifolia*. Thus, while gains may be realized in psyllid resistance of *L. diversifolia* and *L. pallida* relative to *L. leucocephala*, losses in livestock productivity resulting from their use in feeding systems must be anticipated. This is especially important where they are incorporated into concentrate rations as substitutes for expensive commercial protein sources. Other studies have shown that feeding of *A. angustissima* and *C. calothyrsus* as hay has problems of low digestibility and rumen degradability (Ahn *et al.*, 1989; Palmer and Schlick, 1992; Palmer *et al.*, 1994; Dzwowela *et al.*, 1995). In the present study the variability found in chemical composition parameters and IVDMD is interesting. Some of the provenances appear to have dry matter digestibility values that are lower than the generally published data of below 40% for the leaf giving options for improvement through selection. In both species, digestibility of the fodder appears to be strongly influenced by the fibre constituents (ADF and NDF) and the concentration of insoluble proanthocyanidins. Regression analyses in our study have established a high negative relationship, in which the IVDMD (%) relationship with chemical composition parameters in both *Acacia* and *Calliandra* provenances are defined by the regression equations:

$$\text{IVDMD \%} = -0.340 (\text{DM \%}) - 0.941 (\text{NDF \%}) - 0.949 (\text{ADF\%}) + 0.623 (\text{ins. (Acacia) PAS}) + 0.160 (\text{sol. PAS.})$$

$$r^2 = 0.94 \text{ ***}$$

and;

$$\text{IVDMD (\%)} = -0.141 (\text{DM \%}) - 0.928 (\text{NDF \%}) - 0.935 (\text{ADF \%}) + 0.576 (\text{ins. (calliandra) PAS}) + 0.739 (\text{sol. PAS.})$$

$$r^2 = 0.90 \text{ ***}$$

The effect of tannins in fodder legumes on intake of cereal crop residues is important for feeding ruminants in the subhumid Southern Africa because these fodder legume trees can be used as protein supplements (Reed *et al.*, 1990). Legume trees with high content of PAS are associated with low intakes of crop residue whereas those that contain low to moderate levels of PAS are associated with high intakes (Woodward and Reed, 1989). Low total intake and low true digestibility of protein in turn, affect rate of gain for growing animals. In our studies the use of *L. esculenta*, *L. diversifolia* and *L. leucocephala* as supplements has resulted in different utilization efficiencies with *L. diversifolia* producing lower animal performance

than the other two species. From this analysis it is evident that the chemical composition of browse species will affect their potential use in feeding systems. Future research emphasis will be directed to fitting the most promising species and provenances into livestock feed delivery in smallholder production systems.

ACKNOWLEDGEMENTS

We thank the Canadian International Development Agency (CIDA) for funding to the SADC-ICRAF Agroforestry Project, Zimbabwe through which this research was done.

REFERENCES

- Ahn, J.H., B.M. Robertson, R. Elliot, R.C. Gutteridge, and C.W. Ford. 1989. Quality assessment of tropical browse legume: tannin content and protein degradability. *Animal Feed Science and Technology* **27**: 147-156.
- Association of Official Analytical Chemists. 1990. Official Methods of Analyses, 15th Edition. K. Helrich (ed.) AOAC, Washington, DC.
- Dzwowela, B.H. 1993. Advances in forage legumes: sub-Saharan African Perspective pp. 2115-2119. In: Proceedings of the XVII International Grassland Congress, Palmerston North, New Zealand and Rockhampton, Australia.
- Dzwowela, B.H., L. Hove and P.L. Mafongoya. 1995. Effect of drying method on chemical composition and *in vitro* digestibility of multipurpose tree and shrub fodders. *Tropical Grasslands* **29**: 263-269.
- Ndlovu, L.R., L. Mlambo, and B.H. Dzwowela. 1996. Chemical composition, phenolic content and gas production constants of forage of psyllid resistant *Leucaena* species grown in Zimbabwe. *Tropical Grasslands* (submitted).
- Orskov, E.R., F.D. Deb Hovell, and F. Mould. 1980. The use of the nylon bag technique for the evaluation of feedstuffs. *Tropical Animal Production* **5**:195-213
- Palmer, B. and A.C. Schlick. 1992. The effect of drying on the intake and rate of digestion of the shrub legume *Calliandra calothyrsus*. *Tropical Grasslands* **26**: 89-93.
- Palmer, B., D.J. Macqueen and R.A. Bray. 1994. Opportunities and Limitations in *Calliandra* pp 29-34. In: H.M. Shelton, C.M. Piggan and J.L. Brewbaker (eds) proceedings of a workshop. ACIAR Proceedings No. 57, 241 pp.
- Reed, J.D., P.J. Horvath, M.S. Allen and P.J. Van Soest. 1985. Gravimetric determination of soluble phenolics including tannins from leaves by precipitation with trivalent ytterbium. *Journal Science Food Agriculture* **36**:255.
- Robertson, J.B. and P.J. Van Soest. 1981. The detergent system of analysis and its applications to human foods. In: W.P.T. James, and O. Theader, (eds). *The Analysis of Dietary Fibre in Foods*, pp 123-158 (Manel, Dekker, New York)
- Tilley, J.M.A. and R.A. Terry. 1963. A two-stage technique for *in vitro* digestion of forage crops. *Journal of the British Grasslands Society* **18**:104-111.
- Woodward, A. and J.D. Reed. 1989. The influence of polyphenolics on the nutritive value of browse: a summary of research conducted at ILCA. *ILCA Bulletin* **35**:2.

Table 1

Chemical composition of tree fodders from *Leucaena* species compared to native grass hay and IVDMD, DM (%) and N (%) lost after washing and incubating *Leucaena* forage for 24 or 48 hours in the rumen of sheep.

Constituent	Grass	<i>Leucaena</i> species			SED
	Hay	LD	LP	LL	
Dry matter DM, g/kg	950	942	944	943	-
Organic matter (g/kg DM)	950	930	980	940	-
ADF (g/kg DM)	516	272	325	281	-
NDF (g/kg DM)	772	393	459	371	-
N (g/kg DM)	4.3	26.6	18.2	25.1	-
NDI N (%)	74.4	56.4	54.9	47.8	-
Insoluble PAS (ABU. at _{550nm} /g NDF)	1.2	12.8	10.4	8.0	-
IVDMD %	-	48.0	47.8	52.2	-
Washing loss (readily soluble)	DM	19.5	19.9	21.5	1.3
	N	3.7 ^a	10.2 ^b	8.4 ^b	1.4
Rumen degradability after 24 hours	DM	38.1 ^a	52.0 ^c	41.3 ^b	1.5
	N	13.9 ^a	48.4 ^c	29.0 ^b	2.5
Rumen degradability after 48 hours	DM	39.5 ^a	56.0 ^c	48.1 ^b	0.9
	N	19.0 ^a	57.3 ^c	36.9 ^b	1.1

Figures in the row followed by the same letter are not statistically different at P<0.05.

Where NDI N = Neutral detergent insoluble nitrogen

Table 2Chemical composition differences of fodders from *Leucaena* species and provenances

Species Provenances	ID Nos	DM (%)	NDF (%)	Insoluble PAS ^a	CP (%)
<i>L. esculenta</i> subsp. <i>paniculata</i>	OFI 52/87	89.5	36.5	22.5	22.4
<i>L. esculenta</i> subsp. <i>esculenta</i>	OFI 47/87	90.9	40.0	38.2	23.8
<i>L. diversifolia</i> subsp. <i>diversifolia</i>	OFI 45/87	91.2	41.9	29.5	17.0
<i>L. diversifolia</i> subsp. <i>diversifolia</i>	OFI 46/87	90.2	29.5	26.0	19.4
<i>L. diversifolia</i> subsp. <i>stenocarpa</i>	OFI 53/88	92.0	36.3	48.2	17.6
<i>L. diversifolia</i> subsp. <i>stenocarpa</i>	OFI 35/88	91.2	38.5	51.6	17.4
<i>L. leucocephala</i> cv. Cunningham	-	91.2	34.6	15.9	27.6
<i>L. pallida</i>	CPI 58980 offspring	90.1	40.6	63.6	19.5
<i>L. leucocephala</i> x <i>L. diversifolia</i>	ILCA 15090 (Parent=K743)	90.9	39.3	40.2	22.6
<i>L. leucocephala</i> x <i>L. diversifolia</i>	ILCA 15009	90.9	35.1	25.1	19.6
<i>L. shannonii</i> subsp. <i>magnifica</i>	OFI 58/88	92.2	43.6	3.6	24.5
<i>L. shannonii</i> subsp. <i>magnifica</i>	OFI 19/84	91.4	49.3	5.6	25.1
<i>L. pulverulenta</i>	OFI 83/87	90.9	38.9	27.9	22.6
<i>L. salvadorensis</i>	OFI 34/87	93.0	40.0	18.4	25.6
Maize stover	R215	91.5	87.9	nil	3.4
Means (<i>Leucaena</i>)		91.1	38.9	29.7	21.8
	SD±	±0.89	±4.6	±17.2	±3.4

^a (ABU at _{550nm}/gNDF)